



Antileishmanial activity of *Annona* species (Annonaceae)

Heliton Patrick Cordovil Brígido¹ · Juliana Correa-Barbosa¹ · João Victor da Silva-Silva² · Erica Vanessa Souza Costa¹ · Sandro Percário³ · Maria Fâni Dolabela¹

Received: 27 April 2020 / Accepted: 13 August 2020
© Springer Nature Switzerland AG 2020

Abstract

Annona species are widely used in traditional medicine against leishmaniasis. In vitro studies have confirmed their antileishmanial activity. Objective: review the antileishmanial activity of *Annona* species. Results: This article provides a review *Annona* species activity against leishmaniasis, in which it suggests that extracts of *A. mucosa* were active against promastigotes and amastigotes of *L. amazonensis*. Moreover, extracts of *A. crassiflora* were active only against promastigotes of *L. donovani*, whereas the extract, alkaloid fraction and liriiodenine of *A. foetida* were active against promastigotes of *L. braziliensis* and *L. guyanensis*. Liriiodenine was also very active against *L. amazonensis*. Furthermore, extracts and fractions from stems of *A. muricata* were active against *Leishmania* sp. This activity may be related to the presence of acetogenins, since fractionation contributed to increase activity. The fractionation of *A. purpurea* extract contributed to antileishmanial activity, and resulted in a fraction with high selectivity. Such activity may be related to alkaloids or acetogenins. Conclusions: In this review article it is suggested that *Annona* species are promising as leishmanicide and this activity may be related to acetogenins and alkaloids.

Keywords *Annona* · Leishmaniasis · Alkaloid and acetogenin

1 Introduction

Leishmaniasis is caused by parasites belonging to the Trypanosomatidae family and *Leishmania* genus. There are three main forms of leishmaniasis—visceral, cutaneous and mucocutaneous [1]. Most cases of cutaneous leishmaniasis occur in Afghanistan, Algeria, Brazil, Colombia, Islamic Republic of Iran, Pakistan, Peru, Saudi Arabia and Syrian Arab Republic [1].

Leishmaniasis treatment is performed with pentavalent antimonials (sodium stibogluconate and meglumine antimoniate), amphotericin B and pentamidine. However, the use of these agents is questionable due to the variability in their efficacy among *Leishmania* species, high cost, need for parenteral administration, and high toxicity [2, 3].

The search for alternative therapies is very important, and medicinal plants are a source of bioactive molecules [4].

Annona species are used in traditional medicine to treat leishmaniasis. In vitro, antileishmanial studies of extracts validated the popular use [5–7]. Some studies about antileishmanial activity of *Annona* attribute its activity to alkaloids [8, 9] and acetogenins [10].


Several alkaloids were active against *Leishmania*, among these are coronaridine (Fig. 1a), 18-methoxycoronaridine (Fig. 1b) [11], O-methylarmepavine (Fig. 1c) [10], liriiodenine (Fig. 1d) [12]. Moreover, acetogenins annonacinnone (Fig. 1e) and corossolone (Fig. 1f) were also promising as leishmanicide [10].

Previous studies on species of the genus *Annona* present biological investigations on promastigote and amastigote forms of *Leishmania*, for example, from seeds of *A.*

✉ Maria Fâni Dolabela, fanidolabela20@gmail.com | ¹Pharmaceutical Science Post-Graduation Program, Federal University of Pará, Belém, PA, Brazil. ²Laboratory of Immunomodulation and Protozoology, Oswaldo Cruz Institute, Oswaldo Cruz Foundation, Rio de Janeiro, Brazil. ³Oxidative Stress Research Laboratory, Institute of Biological Sciences, Federal University of Pará, Belém, PA, Brazil.



Fig. 1 Compounds isolated from species of *Annona*. Legend: **(a)** Coronaridine, **(b)** 18-Methoxycoronaridine, **(c)** O-methylarmepavine, **(d)** Liriodenine, **(e)** Corossolone, **(f)** Annonacinone



squamosa a trihydroxylated acetogenin with two tetrahydrofuran rings and α , β -unsaturated lactonic ring of 37 carbon atoms endowed with antihelmintic and antiprotozoal properties was isolated. This substance showed leishmanicidal action against promastigotes and amastigotes of *L. chagasi* [10]. The volatile oil of *A. foetida* was active against promastigotes of four different species of *Leishmania*, having been more active in *L. guyanensis* (IC_{50} : 4.1 $\mu\text{g}/\text{mL}$) [9].

The alkaloids fraction from leaves of *A. coriacea* revealed activity against promastigotes of *L. chagasi*, with an IC_{50} of 41.6 $\mu\text{g}/\text{mL}$. In amastigote forms, they caused death to 27.2% of the parasites, at a concentration of 20 $\mu\text{g}/\text{mL}$ [8]. The alkaloid fraction of *A. foetida*, showed activity against the promastigotes of *L. braziliensis* and *L. guyanensis* [12].

In this context, to validate the popular use on leishmaniasis, the analysis of several studies of *Annona* species was performed, focused on their antileishmanial activity.

2 Materials and methods

A survey was performed with the selection of scientific articles available at CAPES, PUBMED and GOOGLE SCHOLAR DATABASE; the year of publication was not limited. The search was carried out in January 2020 and only articles in Portuguese, English and Spanish, which presented leishmanicidal activity (inhibitory concentration 50%— IC_{50}) were considered. The following exclusion criteria were adopted: articles in other languages, those that did not address the proposed theme, in duplicate, and articles that could not be accessed in full.

For the search, descriptors related to the theme were used in an associated way: *Leishmania* and *Annona*. The preliminary result included 1796 papers for screening (CAPES = 21, PUBMED = 15 and GOOGLE SCHOLAR

DATABASE = 1760), using as criterion articles with titles that fit the theme, summary compatible with this study proposal, and the exclusion of duplicate occurrences. Four articles from CAPES and twelve from PUBMED were eliminated due to duplication. Regarding GOOGLE SCHOLAR DATABASE, most of the eliminated articles had inappropriate title or duplication (1757). When the title and abstract were analyzed, 14 papers were included. Figure 2 shows the screening identification procedure included for analysis.

Two reviewers selected, independently, studies based on their title and abstract, those considered potentially relevant were obtained for complete analysis. Any discrepancies were solved by consensus and a third reviewer was consulted to ensure compliance with the inclusion criteria. At the end, 13 articles were selected for discussion and inclusion in this analysis (Fig. 2).




Fig. 2 Flowchart of article eligibility

Regarding data analysis, it was done in two stages: in the first, a table with the following data was used: plant material, type of extract, fractions and isolated substances, IC_{50} and the cytotoxic concentration (CC_{50}). The results are summarized in tables and anti-*Leishmania* activity was assessed using the following criteria: $IC_{50} \leq 100 \mu\text{g/mL}$ active, IC_{50} between 101 and 200 $\mu\text{g/mL}$ moderately active and $IC_{50} \geq 200 \mu\text{g/mL}$ inactive [13]. Cytotoxicity results were assessed by the following criteria: $CC_{50} \leq 100 \mu\text{g/mL}$ cytotoxic, CC_{50} between 101 and 500 $\mu\text{g/mL}$ moderately cytotoxic, and $CC_{50} \geq 500 \mu\text{g/mL}$ non-cytotoxic [13].

3 Results

In order to identify whether *Annona* extracts presented antileishmanial activity, an extensive literature review was done, and the results are highlighted in Table 1. Studies about antileishmanial activity for these species are scarce, with only their anti-promastigote activity being evaluated (Table 1).

The ethanol extracts obtained from root barks, stem barks, and stem wood of *A. crassiflora* were active against promastigotes of *L. donovani* (IC_{50} : $3.7 \pm 0.3 \mu\text{g/mL}$, $12.4 \pm 0.3 \mu\text{g/mL}$, and $8.3 \pm 0.8 \mu\text{g/mL}$, respectively; Table 1) [14]. The total alkaloids obtained from the ethanol extract of *A. crassiflora* leaves were active against *L. chagasi* (IC_{50} : 24.9 ± 0.8 ; Table 1) [8]. The essential oil of *A. foetida* was active against promastigotes of *L. guyanensis* (IC_{50} : $4.1 \pm 0.1 \mu\text{g/mL}$), *L. braziliensis* (IC_{50} : $9.9 \pm 1.2 \mu\text{g/mL}$), *L. amazonensis* (IC_{50} : $16.2 \pm 1.9 \mu\text{g/mL}$), and *L. chagasi* (IC_{50} : $27.2 \pm 6.2 \mu\text{g/mL}$), however, it was cytotoxic in peritoneal macrophages BALB/c (PMBC; CC_{50} : 5.67; Table 1) [9]. The hexane extract from of *A. foetida* was active against promastigotes of *L. guyanensis* (IC_{50} : $42.7 \pm 5.4 \mu\text{g/mL}$). The dichloromethane extract was active against promastigotes of *L. guyanensis* (IC_{50} : $2.7 \pm 0.4 \mu\text{g/mL}$) and *L. amazonensis* (IC_{50} : $23.0 \pm 0.6 \mu\text{g/mL}$). The dichloromethane extract fractionation yielded the alkaloid fraction. However, this fraction showed lower activity against promastigotes of *L. guyanensis* (IC_{50} : $10.3 \pm 0.9 \mu\text{g/mL}$) and *L. amazonensis* (IC_{50} : $18.3 \pm 2.5 \mu\text{g/mL}$). Nevertheless, the methanol extract of *A. foetida* was active against *L. guyanensis* (IC_{50} : $23.6 \pm 3.1 \mu\text{g/mL}$) and *L. amazonensis* (IC_{50} : $40.4 \pm 3.2 \mu\text{g/mL}$), and fractionation contributed to the activity (*L. guyanensis* IC_{50} : $9.1 \pm 0.8 \mu\text{g/mL}$ and *L. amazonensis* IC_{50} : $24.3 \pm 1.9 \mu\text{g/mL}$; Table 1) [12].

The antileishmanial activity of alkaloids and acetogenins isolated from *A. foetida* were evaluated. The alkaloid liriodenine was more promising against *L. guyanensis* (IC_{50} : $21,0.5 \pm 0.4 \mu\text{g/mL}$ and $55.92 \pm 3.55 \mu\text{g/mL}$) [9, 15] and PH8- *L. amazonensis* (IC_{50} : $1.43 \pm 0.58 \mu\text{g/mL}$). Liriodenine showed high toxicity for BALB/c mice peritoneal

macrophages (CC_{50} : $19.11 \pm 1.06 \mu\text{g/mL}$) and low selectivity in *L. guyanensis* (SI: 0.34; Table 1) [15].

Different extracts obtained from *Annona mucosa* were tested on *L. donovani*, *L. amazonensis*, and *L. braziliensis*. Most of the extracts showed activity against promastigotes of *L. amazonensis* (PH8, IC_{50} : 9.3–46.5 $\mu\text{g/mL}$), although it was not observed significant reduction in the rate of macrophages infection by amastigotes (30%). When cytotoxicity and anti-promastigote activity are related, a low selectivity index is observed (SI: 0.9–6; Table 1) [15].

From *Annona mucosa*, the alkaloids oxoaporphine, atherospermidina and liriodenine were isolated. Liriodenine was active against promastigotes of *L. amazonensis* (IC_{50} : $1.43 \pm 0.58 \mu\text{g/mL}$) and amastigotes forms of *L. amazonensis* [15]. High selectivity was observed (SI: 13.36), especially for liriodenine (SI: 13.37; Table 1).

The anti-promastigote activity of *A. muricata* was extensively evaluated, yielding inactive (hexane and methanol extracts from pericarp: $IC_{50} > 1000 \mu\text{g/mL}$), moderately active (hexane and methanol extracts from leaves: $IC_{50} > 100 \mu\text{g/mL}$), and active extracts (hexane and methanol extracts from stem and ethyl acetate extracts from pericarp, leaves, and stem: $IC_{50} \leq 100 \mu\text{g/mL}$). Fractionation of *A. muricata* extracts led to the isolation of acetogenins, which were more active than the extracts against promastigotes (Table 1) [10, 16].

Scoparone, corosolone, and annonacinone isolated from *A. muricata* were active against promastigotes of *L. donovanni*, *L. mexicana*, and *L. major* [10, 16]. Annonacinone displayed the major activity against those three *Leishmania* species (IC_{50} : 6.72–8.00 $\mu\text{g/mL}$) [16]. Annonacinone (IC_{50} : 37.6 $\mu\text{g/mL}$) and corosolone (IC_{50} : 25.9 $\mu\text{g/mL}$) showed less activity against promastigotes of *L. chagasi*. These substances were also tested against amastigotes of *L. chagasi*, being annonacinone (IC_{50} : 13.5 $\mu\text{g/mL}$) more active than corosolone (IC_{50} : 28.7 $\mu\text{g/mL}$). Annonacinone also presented higher selectivity (CC_{50} : 59.5 $\mu\text{g/mL}$; SI: 4.4) than corosolone (CC_{50} : 54 $\mu\text{g/mL}$; SI: 1.9; Table 1) [10].

Scoparone presented lower activity than others acetogenins against promastigotes of *L. donovanni* (IC_{50} : $27.51 \pm 0.97 \mu\text{g/mL}$), but presented activity similar to the others compounds against *L. mexicana* (IC_{50} : $9.11 \pm 0.25 \mu\text{g/mL}$) and *L. major* (IC_{50} : $14.37 \pm 0.98 \mu\text{g/mL}$; Table 1) [16].

The alkaloid O-methylarmepavine and the acetogenin C_{37} trihydroxy adjacent bistetrahydrofuran were isolated from *A. squamosa*. Both O-methylarmepavine and trihydroxy adjacent bistetrahydrofuran showed similar inhibitory effects against promastigotes (IC_{50} : 23.3 and 26.4 $\mu\text{g/mL}$) and amastigotes of *L. chagasi* (IC_{50} : 25.4 and 25.3 $\mu\text{g/mL}$). However, this acetogenins showed higher cytotoxicity in RAW 264.7 cells (CC_{50} : 43.5 $\mu\text{g/mL}$) than the alkaloid (CC_{50} : 79.7 $\mu\text{g/mL}$; Table 1) [16].

Table 1 Antileishmanial activity of *Annona* species

Samples	Antileishmanial activity (IC ₅₀ µg/mL ± SD)		<i>Leishmania</i> species and cell	Cytotoxicity (CC ₅₀ µg/mL)	Activity classification	Selectivity index (SI)	References
	Promastigote	Amastigote					
<i>Annona crassiflora</i>							
EE SB	12.4 ± 0.3	ND	<i>L. donovani</i>	ND	Active	ND	Mesquita et al. (2005)
EE SW	8.3 ± 0.8	ND	<i>L. donovani</i>	ND	Active	ND	Mesquita et al. (2005)
EE RB	3.7 ± 0.3	ND	<i>L. donovani</i>	ND	Active	ND	Mesquita et al. (2005)
EE RW	8.7 ± 0.6	ND	<i>L. donovani</i>	ND	Active	ND	Mesquita et al. (2005)
Total alkaloids of leaves	24.9 ± 0.8	ND	<i>L. chagasi</i>	ND	Active	ND	Tempone et al. (2005)
<i>Annona foetida</i>							
HE bark	> 160 42.7 ± 5.4	ND	<i>L. braziliensis</i> <i>L. guyanensis</i>	ND	Moderate or inactive Active	ND	Costa et al. (2006)
DCE bark	23.0 ± 0.6 2.7 ± 0.4	ND	<i>L. braziliensis</i> <i>L. guyanensis</i>	ND	Active Active	ND	Costa et al. (2006)
Alkaloid fraction (DCE)	18.3 ± 2.5 10.3 ± 0.9	ND	<i>L. braziliensis</i> <i>L. guyanensis</i>	ND	Active Active	ND	Costa et al. (2006)
ME bark	40.4 ± 3.2 23.6 ± 3.1	ND	<i>L. braziliensis</i> <i>L. guyanensis</i>	ND	Active Active	ND	Costa et al. (2006)
Alkaloid fraction (ME)	24.3 ± 1.9 9.1 ± 0.8	ND	<i>L. braziliensis</i> <i>L. guyanensis</i>	ND	Active Active	ND	Costa et al. (2006)
Alkaloid N-hydroxyannonomontine	252.7 ± 2.2 437.5 ± 2.5	ND	<i>L. braziliensis</i> <i>L. guyanensis</i>	ND	Inactive Inactive	ND	Costa et al. (2006)
Alkaloid O-methylmoschatolin	320.8 ± 3.1 103.7 ± 3.4	ND	<i>L. braziliensis</i> <i>L. guyanensis</i>	ND	Inactive Moderate active	ND	Costa et al. (2006)
Alkaloid liriodenine	58.5 ± 1.8 21.5 ± 0.4	ND	<i>L. braziliensis</i> <i>L. guyanensis</i>	ND	Active Active	ND	Costa et al. (2006)
Alkaloid annonmontine	34.8 ± 1.5 > 613.0	ND	<i>L. braziliensis</i> <i>L. guyanensis</i>	ND	Active Inactive	ND	Costa et al. (2006)
Essential Oil	16.2 ± 1.9 9.9 ± 1.2 27.2 ± 6.2 4.1 ± 0.1	ND	<i>L. amazonensis</i> <i>L. braziliensis</i> <i>L. chagasi</i> <i>L. guyanensis</i>		Active Active Active Active	0.35 0.57 0.20 1.38	Costa et al. (2009)
			PMBC	5.67	Cytotoxic		
<i>Annona mucosa</i>							
HE of leaves	24.24 ± 1.51 65.27 ± 1.20	ND	PH8 M2903		Active Active	2.58 0.95	Lima et al. (2012)
			PMBC	62.63 ± 4.10	Cytotoxic		
HE of seeds	44.22 ± 5.64 170.15 ± 1.46	ND	PH8 M2903		Active Moderate active	5.93 1.54	Lima et al. (2012)
			PMBC	262.33 ± 5.81	Moderate cytotoxicity		
DCE of leaves	9.32 ± 0.56 27.42 ± 5.42	ND	PH8 M2903		Active Active	2.58 0.87	Lima et al. (2012)
			PMBC	24.07 ± 0.72	Cytotoxic		

Table 1 (continued)

Samples	Antileishmanial activity (IC ₅₀ µg/mL ± SD)		<i>Leishmania</i> species and cell	Cytotoxicity (CC ₅₀ µg/mL)	Activity classification	Selectivity index (SI)	References
	Promastigote	Amastigote					
ME of leaves	28.32 ± 1.15	ND	PH8		Active	1.03	Lima et al. (2012)
	44.74 ± 5.89		M2903		Active	0.65	
			PMBC	29.41 ± 0.89	Cytotoxic		
ME of seeds	46.54 ± 4.95	ND	PH8		Active	2.98	Lima et al. (2012)
	133.17 ± 5.41		M2903		Moderate active	1.04	
			PMBC	139.0 ± 3.13	Moderate cytotoxicity		
Alkaloid lirioidenine	1.43 ± 0.58	ND	PH8		Active	13.36	Lima et al. (2012)
	55.92 ± 3.55		M2903		Active	0.34	
			PMBC	19.11 ± 1.06	Cytotoxic		
HE of leaves	100.0	ND	PH8	ND	Active	ND	Osorio et al. (2007)
	> 100.0		M2903		Moderate or inactive		
	> 100.0		PP75		Moderate or inactive		
HE of stem	98.6	ND	PH8	ND	Active	ND	Osorio et al. (2007)
	76.3		M2903		Active		
	83.1		PP75		Active		
EAE of leaves	25.0	ND	PH8		Active	0.312	Osorio et al. (2007)
	25.0		M2903		Active	0.312	
	25.0		PP75		Active	0.312	
EAE of stem	63.2	ND	PH8	7.8 ± 0.3	Cytotoxic		Osorio et al. (2007)
	63.2		M2903	ND	Active	ND	
	63.2		PP75		Active		
ME of leaves	> 100.0	ND	PH8	NDND	Moderate or inactive	ND	Osorio et al. (2007)
	> 100.0		M2903		Moderate or inactive		
	> 100.0		PP75		Moderate or inactive		
ME of stem	98.6	ND	PH8	ND	Active	ND	Osorio et al. (2007)
	98.6		M2903		Active		
	98.6		PP75		Active		
Acetogenin corosolone	25.9	28.7	<i>Leishmania chagasi</i>		Active	2.08	Villa-Nova et al. (2011)
Acetogenin annonacinone	37.6	13.5	RAW 264.7	54.0	Cytotoxic		Villa-Nova et al. (2011)
			<i>Leishmania chagasi</i>	59.5	Active	1.58	
Acetogenin scoparone	27.51 ± 0.97	ND	RAW 264.7		Cytotoxic		Villa-Nova et al. (2013)
	9.11 ± 0.25		<i>L. donovani</i>	ND	Active	ND	
	14.37 ± 0.98		<i>L. mexicana</i> <i>L. major</i>		Active Active		

Table 1 (continued)

Samples	Antileishmanial activity (IC ₅₀ µg/mL ± SD)		<i>Leishmania</i> species and cell	Cytotoxicity (CC ₅₀ µg/mL)	Activity classification	Selectivity index (SI)	References
	Promastigote	Amastigote					
Acetogenin corosolone	18.73 ± 0.82	ND	<i>L. donovani</i>	ND	Active	ND	Villa-Nova et al. (2013)
	18.64 ± 0.79		<i>L. mexicana</i>		Active		
	16.14 ± 1.13		<i>L. major</i>		Active		
Acetogenin annonacinone	7.66 ± 0.77	ND	<i>L. donovani</i>		Active	ND	Villa-Nova et al. (2013)
	8.00 ± 1		<i>L. mexicana</i>		Active		
	6.72 ± 0.37		<i>L. major</i> <i>Annona squamosa</i>		Active		
Alkaloid O-methyl-armepavine	23.3	25.4	<i>L. chagasi</i>		Active	3.42	Villa-Nova et al. (2011)
Acetogenin	26.4	25.3	RAW 264.7	79.7	Cytotoxic		Villa-Nova et al. (2011)
			<i>L. chagasi</i>	43.5	Active	1.64	
C ₃₇ trihydroxy adjacent bistetrahydrofuran			RAW 264.7		Cytotoxic		
HAE of leaves	37.8 ± 0.1	ND	<i>Annona glabra</i> <i>L. amazonensis</i> <i>Annona senegalensis</i>	ND	Active		García et al. (2012)
EE of leaves	10.8	ND	<i>L. donovani</i> JURKAT	273.49	Active Moderate cytotoxicity	25.32	Ohashi et al. (2018)
EE of stem cutting	27.8	ND	<i>L. donovani</i> JURKAT	127.95	Active Moderate cytotoxicity	4.60	Ohashi et al. (2018)
ME of bark	113.24 ± 1.2	ND	<i>Annona purpurea</i> <i>L. donovani</i>		Moderate active	0.0008	Camacho et al. (2003)
			KB	0.0098	Cytotoxic		
H ₂ O of bark	289 ± 3.70	ND	<i>L. donovani</i>		Inactive	ND	Camacho et al. (2003)
			KB	> 500	Non cytotoxic		
ME of seed	28.57 ± 1.78	ND	<i>L. donovani</i>	7.81 ± 1.45	Active	0.27	Camacho et al. (2003)
			KB		Cytotoxic		
H ₂ O of seed	179.9 ± 4.1	ND	<i>L. donovani</i>	96.7 ± 3.9	Moderate active	0.53	Camacho et al. (2003)
			KB		Cytotoxic		
E ₂ fraction of HAE leaves	0.961	ND	<i>L. panamensis</i>	124.02	Active	129.05	Cárdenas et al. (2005)
			U937		Moderate cytotoxicity		
Extract	175	ND	<i>Annona cornifolia</i> <i>L. amazonensis</i>		Moderate active	0.196	De Toledo et al. (2011)
			VERO	34.33	Cytotoxic		

HAE: hydroalcoholic extract; HE: Hexane extract; EAE: ethyl acetate extract; ME: methanol extract; CE: chloroform extract; DCE: Dichloromethane extract; SB: stem bark; SW: stem wood; RB: root bark; KB: Human nasopharyngeal KB cells; RAW 264.7: Murine macrophage cells; PMBC: Peritoneal macrophage BALB-C; U937: Human monocytic cell lineage; PH8: *Leishmania amazonensis*; PP75: *Leishmania donovani*; M2903: *Leishmania braziliensis*; ND: Not determined. JURKAT: human acute T-cell leukemia cells

The hydroalcoholic leaf extract of *A. glabra* was active against *L. amazonensis* promastigotes (IC_{50} : $37.8 \pm 0.1 \mu\text{g/mL}$) [17]. The extracts obtained from the leaves and branches of *A. senegalensis* also showed activity against another strain of Leishmania (*L. donovani*; IC_{50} 10.8 $\mu\text{g/mL}$ and 27.8 $\mu\text{g/mL}$ respectively) [18]. However, the two samples were moderately cytotoxic for human T-cell of acute leukemia (JURKAT; CC_{50} : 273.49 $\mu\text{g/mL}$ and 127.95 $\mu\text{g/mL}$ respectively) [18] and the leaf extract showed best selectivity index (SI: 25.3; Table 1).

The methanol and aqueous extracts from *A. purpurea* bark and seed were evaluated against *L. donovani* promastigotes. The methanol extract from the seed showed better activity (IC_{50} : $28.57 \pm 1.78 \mu\text{g/mL}$) but was cytotoxic in nasopharyngeal cells. (KB; CC_{50} : 7.81 ± 1.45 ; IS: 0.27). However, the aqueous seed extract presented the best selectivity index (IC_{50} : $179.9 \pm 4.1 \mu\text{g/mL}$; CC_{50} : $96.7 \pm 3.9 \mu\text{g/mL}$; SI: 0.53; Table 1) [19]. The fraction E2 obtained from the hydroalcoholic leaf extract of *A. purpurea* was active against *L. panamensis* (IC_{50} : 0.961) and moderately cytotoxic in human monocyte cells, presenting a high selectivity index (U937; CC_{50} : 124.02 $\mu\text{g/mL}$; SI: 129.05; Table 1) [20].

The extract from *A. cornifolia* was moderately active in *L. amazonensis* (IC_{50} : 175 $\mu\text{g/mL}$) and cytotoxic in African green monkey kidney cells (VERO; CC_{50} : 34.33 $\mu\text{g/mL}$; SI: 0.196; Table 1) [21]. Other extracts from branches and leaves of *A. muricata* were active against promastigotes of *L. amazonensis*, *L. braziliensis* and *L. donovani* (Table 1) [22].

The ethyl acetate extracts from both the branch and the leaves showed better results in the 3 *Leishmania* species (branch: IC_{50} : 63.2 $\mu\text{g/mL}$; leaves: 25 $\mu\text{g/mL}$; Table 1) [22].

4 Discussion

Different species belonging to the *Annona* genus have popular use for leishmaniasis or wounds treatment [7, 10, 12]. However, there is the following question: this activity is related to the presence of acetogenins, or their alkaloids, or it is related to the synergism between these metabolites.

Previous study demonstrated anomontine alkaloid was the most active against *L. amazonensis* [12]. Alkaloids can interfere with tubulin polymerization or stabilize the DNA topoisomerase complex [23]. The biological activity of liriodenine can be attributed to the intercalation between neighboring base pairs of the DNA double helix [24, 25].

The oxo function induces cytotoxicity on precursor incorporation into DNA [25, 26]. Indeed, acetogenins binds to complex I of the mitochondrial electron

transport chain [27, 28]. In addition, it inhibits ubiquinone-bound NADH oxidase [29–31]. Due to the differences in mechanisms of action, it is believed that there may be synergy to leishmanicide activity.

In the present review, although acetogenins were very active against promastigotes of different species of *Leishmania*, with an IC_{50} of less than 30 $\mu\text{g/mL}$ [10, 16] corosolone and annonacinone were still active against the amastigote form [10]. In contrast, structural changes seem to interfere in the antileishmanial activity of alkaloids and this activity may be more pronounced depending on the parasite species [10, 12, 15].

If only antiparasitic activity is considered, this may suggest that acetogenins are more promising than alkaloids as leishmanicide. However, you must consider the toxicological aspects and the selectivity index (SI).

Unfortunately, most of the acetogenins isolated from *Annona* lack cytotoxicity studies, and preliminary results suggest that corosolone (SI = 2.08) and annonacinone (SI = 1.58) have a selectivity index of less than 10 [10]. However, additional genotoxicity, mutagenicity and in vivo studies need to be carried out to determine the safety of its use as a leishmanicide.

Another issue that needs to be analyzed is whether the fractionation contributes to antileishmanial activity. In the case of *A. muricata*, fractionation contributed significantly to biological activity, it also seems to contribute to selectivity [7, 22].

Nevertheless, for *A. crassiflora*, some extracts showed better antileishmanial potential than isolated substances [8, 14]. For *A. foetida*, alkaloid fractions appear to be promising for leishmaniasis treatment (Table 1) [12].

The fractionation of *A. purpurea* extract contributed to antileishmanial activity and high selectivity index, with the E2 fraction being a strong candidate for anti-leishmania drug, since this showed better anti-leishmania activity (IC_{50} : 0.961 $\mu\text{g/mL}$), moderate cytotoxicity, resulting in a selectivity index > 10 among the extracts, fractions and isolated substances evaluated from *Annona* species (IS: 129.05) [20].

5 Conclusion

Annona species are promising for the treatment of leishmaniasis, bearing in mind the importance of extract fractionation and alkaloids or acetogenins isolation, since many alkaloids and acetogenins have antiparasitic activity, which can be important tools against parasites drug resistant.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

References

1. Who Leishmaniasis Who (2018) <https://www.who.int/leishmaniasis/en/>. Accessed January 25, 2020
2. Croft SL, Coombs GH (2003) Leishmaniasis—current chemotherapy and recent advances in the search for novel drugs. *Trends Parasitol* 19(11):502–508
3. Oliveira RA, Lima EO, Vieira WL et al (2006) Estudo da interferência de óleos essenciais sobre a atividade de alguns antibióticos usados na clínica. *Rev Bras Farmacogn* 16(1):77–82
4. da Rocha FAG, de Araújo LSG, Lima TGD et al (2013) Características do comércio informal de plantas medicinais no município de Lagoa Nova/RN. *HOLOS* 5:264–281
5. Sahpaz S, Bories CH, Loiseau PM, Cortes D, Hocquemiller R, Laurens A, Cave A (1994) Cytotoxic and antiparasitic activity from *Annona senegalensis* seeds. *Planta Med* 60(06):538–540
6. Saez J, Granados H, Torres B, Velez ID, Munoz D (1998) Leishmanicidal activity of *Annona* aff. *spraguei* seeds. *Fitoterapia (Milano)* 69(5):478–479
7. Jaramillo MC, Arango GJ, González MC, Robledo SM, Velez ID (2000) Cytotoxicity and antileishmanial activity of *Annona muricata* pericarp. *Fitoterapia* 71(2):183–186
8. Tempone AG, Borborema ST, de Andrade Jr HF et al (2005) Antiprotozoal activity of Brazilian plant extracts from isoquinoline alkaloid-producing families. *Phytomedicine* 12(5):382–390
9. Costa EV, Pinheiro MLB, Silva JRDA et al (2009) Antimicrobial and antileishmanial activity of essential oil from the leaves of *Annona foetida* (Annonaceae). *Quím Nova* 32(1):78–81
10. Vila-Nova NS, Morais SMD, Falcão MJC et al (2011) Leishmanicidal activity and cytotoxicity of compounds from two Annonaceae species cultivated in Northeastern Brazil. *Rev Soc Bras Med Trop* 44(5):567–571
11. Delorenzi JC, Freire-de-Lima L, Gattass CR, Costa DA, He L, Kuehne ME et al (2002) In vitro activities of Iboga alkaloid congeners coronaridine and 18-methoxycoronaridine against *Leishmania amazonensis*. *Antimicrob Agents Chemother* 46:2111–2115
12. Costa EV, Pinheiro MLB, Xavier CM et al (2006) A pyrimidine- β -carboline and other alkaloids from *Annona foetida* with Antileishmanial Activity. *J Nat Prod* 69(2):292–294
13. da Silva e Silva JV, Brígido CPH, Oliveira de Albuquerque KC, et al (2019) Flavopereirine—An alkaloid derived from *geissospermum vellosii*—Presents leishmanicidal activity in vitro. *Molecules* 24(4):785
14. Mesquita MLD, Desrivot J, Fournet A, Paula JED, Grellier P, Espindola LS (2005) Antileishmanial and trypanocidal activity of Brazilian Cerrado plants. *Mem Inst Oswaldo Cruz* 100(7):783–787
15. De Lima JP, Pinheiro ML, Santos AMG et al (2012) In vitro antileishmanial and cytotoxic activities of *Annona mucosa* (Annonaceae). *Rev Virtual Quim* 4(6):692–702
16. Vila-Nova NS, Morais SMD, Falcão MJC et al (2013) Different susceptibilities of *Leishmania* spp. promastigotes to the *Annona muricata* acetogenins annonacinone and corosolone, and the *Platymiscium floribundum* coumarin scoparone. *Ex parasitol* 133(3):334–338
17. García M, Monzote L, Scull R, Herrera P (2012) Activity of Cuban plants extracts against *Leishmania amazonensis*. *ISRN pharmacol* 2012
18. Ohashi M, Amoa-Bosompem M, Kwofie KD et al (2018) In vitro antiprotozoan activity and mechanisms of action of selected Ghanaian medicinal plants against *Trypanosoma*, *Leishmania*, and *Plasmodium* parasites. *Phytother Res* 32(8):1617–1630
19. Camacho MDR, Phillipson JD, Croft SL, Solis PN, Marshall SJ, Ghazanfar SA (2003) Screening of plant extracts for antiprotozoal and cytotoxic activities. *J Ethnopharmacol* 89(2–3):185–191
20. Cárdenas DL, Lora JA, Márquez RL, Blanco PJ (2005) Actividad leishmanicida de *Annona purpurea*. *Actual Biol* 27(1):35–37
21. De Toledo CE, Britta EA, Ceole LF et al (2011) Antimicrobial and cytotoxic activities of medicinal plants of the Brazilian cerrado, using Brazilian cachaça as extractor liquid. *J Ethnopharmacol* 133(2):420–425
22. Osorio E, Arango GJ, Jiménez N et al (2007) Antiprotozoal and cytotoxic activities in vitro of Colombian Annonaceae. *J ethnopharmacol* 111(3):630–635
23. Cheng X, Wang D, Jiang L, Yang D (2008) DNA topoisomerase I inhibitory alkaloids from *Corydalis saxicola*. *Chem Biodivers* 5(7):1335–1344
24. Woo SH, Reynolds MC, Sun NJ, Cassady JM, Snapka RM (1997) Inhibition of topoisomerase II by liriiodenine. *Biochem Pharmacol* 54(4):467–473
25. Chen CY, Wu HM, Chao WY, Lee CH (2013) Review on pharmacological activities of liriiodenine. *Afr J Pharm Pharmacol* 7:1067–1070
26. Tzeng CC, Wu YC, Su TL, Watanabe KA, Lu ST, Chou TC (1990) Inhibitory effects of isoquinoline-type alkaloids on leukemic cell growth and macromolecule biosynthesis. *Gaoxiong yi xue ke xue za zhi= Kaohsiung J Med Sci* 6(2):58–65
27. Gallardo T, Aragón R, Tormo J, Blázquez MA, Zafra-Polo MC, Cortes D (1998) Acetogenins from *Annona glabra* seeds. *Phytochemistry* 47(5):811–816
28. Guadaño A, Gutiérrez C, de la Peña E, Cortes D, González-Coloma A (2000) Insecticidal and mutagenic evaluation of two annonaceous acetogenins. *J Nat Prod* 63(6):773–776
29. Zeng L, Wu FE, Oberlies NH, McLaughlin JL, Sastrodihadjo S (1996) Five new monotetrahydrofuran ring acetogenins from the leaves of *Annona muricata*. *J Nat Prod* 59(11):1035–1042
30. Gleye C, Raynaud S, Hocquemiller R et al (1998) Muricadienins, muridienins and chatenaytrienins, the early precursors of annonaceous acetogenins. *Phytochemistry* 47(5):749–754
31. Allali FQ, Liu X, McLaughlin JL (1999) Annonaceous acetogenins: recent progress. *J Nat Prod* 62:504–540

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.